

Driving, drug use behaviour and risk perceptions of nightclub attendees in Victoria, Australia.

Author(s)

Degenhardt L, Dillon P, Duff C et al.

Published

Sunday, January 1, 2006 - 12:00

Publisher

International Journal of Drug Policy

Volume

17

Issue

1

Page(s)

41-46

Abstract

Aim: In anticipation of the introduction of roadside saliva testing in Victoria for recent THC and methylamphetamine use, this study examined the prevalence of illicit drug use among nightclub attendees in Melbourne, Victoria; their transport methods; and their drug use and driving histories. **Methods:** In total, 273 persons were interviewed as they entered nightclubs in the inner Melbourne area. Questionnaires addressed drug use, risk perception and driving behaviour. **Results:** Drug use on the night of interview was common, with notable proportions reporting that they had used or intended to use cannabis (22%), ecstasy (18%), speed (13%) and crystal/base methamphetamine (6%). Around one in ten participants reported that on the night of interview, they would either drive or be driven by someone under the influence of alcohol (10%), cannabis (11%) and/or methamphetamine (8%). Seventy percent reported having heard of roadside drug testing and 65% supported it. Forty percent reported that roadside drug testing would change their drug driving behaviour. **Conclusions:** Roadside drug testing in Victoria may have positive impacts upon drug use and driving risk behaviours among a sample of young persons attending nightclubs. Information provision and increasing the transport options for young people will play a part in reducing the number of young

clubbers who use drugs and drive. (PsycINFO Database Record (c) 2009 APA, all rights reserved) (from the journal abstract)

Web link

<http://linkinghub.elsevier.com/retrieve/pii/S0955395905001957>

[View PDF](#)